

METHOD FOR TREATING
CHRONIC MYELOGENOUS LEUKEMIA

Inventor: John Lyons

BACKGROUND OF THE INVENTION

Field of the Invention

The invention relates to methods, compositions, and kits for treating cancer associated with protein tyrosine kinase activity, and more particularly for treating chronic myelogenous leukemia.

Description of Related Art

Chronic Myelogenous Leukemia (CML) is a myeloproliferative disorder of a pluripotent hematopoietic stem cell with a particular cytogenetic abnormality, the Philadelphia chromosome. Faderl et al (1999) Ann. Intern. Med. 131: 207-219. In childhood, it accounts for only 2 to 5 % of all malignant disorders and presents as either of two distinct clinical entities, adult-type CML and juvenile CML. Adult-type CML of childhood is indistinguishable from that seen in older patients. However, juvenile CML is restricted to children and is Philadelphia chromosome negative. Grier and Civin (1998) in (Nathan and Oski, eds) Hematology of Infancy and Childhood, volume 2, 5th ed, W. B. Saunders Company, 34:1286-1459.

CML is a progressive, uniformly fatal disease in untreated patients. It is characterized by three distinct phases: a chronic phase lasting three to five years; an acute or accelerated phase lasting three to six months; and a brief blastic crisis phase. The progression of the disease to blast crisis results in rapid death due to infections, bleeding and leukemic organ infiltration.

Philadelphia chromosome, the characteristic cytogenetic abnormality of CML, results from a reciprocal chromosomal translocation, t(9;22)(q34;q11), in a

hematopoietic stem cell. This translocation produces a fusion gene, termed Bcr-Abl, created by the juxtaposition of the abelson murine leukemia (Abl) protooncogene on chromosome 9 with a portion of the breakpoint cluster region (Bcr) gene on chromosome 22. The Bcr-Abl fusion protein's leukomogenic potential is derived from its constitutively activated tyrosine kinase activity, which causes a perturbation of stem cell function through unclear mechanisms. This activity results in interference with basic cellular processes, such as control of proliferation, adherence, and physiological death. More advanced stages of CML are also characterized by aberrant methylation of multiple genes, including the p15/Ink-4b cell-cycle regulator gene. Cortes et al (1997) *Baillieres Clin Haematol* 10(2):277-90. Aberrations in DNA methylation, whether general or site specific, are common in cancer and have important roles in tumor initiation, progression and resistance. Lubbert et al (2001) *Br J Haematol* 114(2):349-357.

Until recently, standard therapy for chronic phase CML consisted of conventional chemotherapy, interferon-alpha (with or without Ara-C), donor lymphocyte infusions and allogeneic bone marrow transplantation, each offering different risk-benefit trade-offs.

Overall, available data suggests the view that allogeneic bone marrow transplantation offers to eligible patients (children and young adults with an human leukocyte antigen-matched sibling donor) their best prospect for cure. However, bone marrow transplantation has certain limitations i.e., the availability of a suitable donor (10-40 %), the risk of graft-vs.-host disease (8-60%) and a high rate of transplant-related mortality (20-40%).

Long-term follow-up of patients treated in large-scale randomized trials utilizing one or two of the above therapeutic modalities has shown a significant correlation between cytogenetic responses and prolonged survival. Silver et al (1999) *Blood* 94:1517-1536.

Imatinib mesylate is one of the recent therapeutic breakthroughs in the treatment of CML. Imatinib mesylate is a small molecule inhibitor of tyrosine kinase activity that results in a high response rate in CML. In the pivotal Phase II

studies, nearly all patients (88%) with chronic phase CML achieved a complete hematologic response, and nearly half (49%) had a major cytogenetic response. Imatinib mesylate produced remissions in 63% of accelerated phase CML patients and 26% of blast phase patients. A complete cytogenetic response was seen in 30% of chronic phase CML, 14% of accelerated phase CML, and 5% of blast phase CML patients and was maintained for four weeks in 16%, 4%, and 1%, respectively. Novartis, Gleevec package insert T-2001-14 90012401.

Imatinib mesylate was approved by the FDA in May 2001 for the treatment of CML in all phases (after failure of interferon in the chronic phase). However, in blast phase CML, the responses to imatinib mesylate are usually of very short duration, and most patients manifest resistant/refractory disease within six months of therapy. Druker et al (2001) N. Engl. J. Med. 344: 1038-1042. Resistance to imatinib mesylate was associated with reactivation of Bcr-Abl and could be conferred by a single point substitution of threonine for isoleucine in the tyrosine kinase. Gorre et al (2001) Science 293: 2163. Consequently, there exists a need for compositions and methods for treating CML patients who are resistant to imatinib mesylate.

SUMMARY OF THE INVENTION

A method is provided for treating a patient with imatinib mesylate in combination with a DNA methylation inhibitor. The method is preferably directed to a patient that has a degree of resistance to imatinib mesylate, the resistance being mitigated by the administration of the DNA methylation inhibitor. In particular, the method is directed to treating a disease state associated with activity of protein tyrosine kinase such as oncoprotein Bcr-Abl involved in chronic myelogenous leukemia (CML), platelet-derived growth factor (PDGF) receptor involved in prostate cancer and glioblastoma, and c-Kit involved in gastrointestinal stromal tumor (GIST) and small cell lung cancer (SCLC), as well as other types of cancer,

where the combination treatment using imatinib mesylate and a DNA methylation inhibitor is synergistic.

In one embodiment, the method comprises: administering to the patient imatinib mesylate and a DNA methylation inhibitor.

In another embodiment, a method is provided for treating a patient having chronic myelogenous leukemia comprising: administering to a patient having chronic myelogenous leukemia and a degree of resistance to imatinib mesylate, a therapeutically effective amount of a DNA methylation inhibitor which mitigates the imatinib mesylate resistance.

In one variation, the patient has already manifested resistance to imatinib mesylate within 6 months of the treatment with imatinib mesylate as defined by no improvement in the prognosis or worsening of the prognosis.

In another embodiment, a method is provided for treating a patient having chronic myelogenous leukemia comprising: administering to a patient having chronic myelogenous leukemia and manifesting intolerance to imatinib mesylate, a therapeutically effective amount of a DNA methylation inhibitor which mitigates the imatinib mesylate intolerance.

In one variation, the patient has already manifested intolerance to imatinib mesylate within 6 months of the treatment with imatinib mesylate as defined by manifesting a symptom selected from the group consisting of hepatotoxicity, fluid retention syndrome, neutropenia, hemorrhage, dyspepsia, dyspnea, diarrhea, muscle cramps, skin rash, fatigue, headache, nausea, vomiting, and thrombocytopenia.

In another embodiment, a method is provided for treating a patient having chronic myelogenous leukemia comprising: administering to the patient imatinib mesylate and a DNA methylation inhibitor.

In another embodiment, a method is provided for treating a patient having chronic myelogenous leukemia and a degree of resistance to imatinib mesylate comprising: administering to the patient imatinib mesylate and a DNA methylation inhibitor such that the patient's resistance to imatinib mesylate in the absence of the DNA methylation inhibitor is reduced.

In another embodiment, a method is provided for treating a patient having chronic myelogenous leukemia, comprising: administering to a patient in blast phase of chronic myelogenous leukemia a therapeutically effective amount of a DNA methylation inhibitor.

According to any of the above methods for treating chronic myelogenous leukemia, the patient's chronic myelogenous leukemia is optionally staged prior to administration. Staging the patient having chronic myelogenous leukemia optionally includes determining the number of blasts, promyelocytes, basophil, and platelets per liter of peripheral blood or bone marrow.

Also according to any of the above methods for treating chronic myelogenous leukemia, the DNA methylation inhibitor is optionally administered to the patient in the blast, chronic or accelerated phase of chronic myelogenous leukemia. In one variation, the method is performed when the patient in blast phase of chronic myelogenous leukemia has more than 30% blasts in peripheral blood or bone marrow.

Also according to any of the above methods for treating chronic myelogenous leukemia, the DNA methylation inhibitor may be administered by a variety of routes, including but not limited to orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery, subcutaneously, intraadiposally, intraarticularly, or intrathecally.

According to any of the above methods, it is noted that administering imatinib mesylate and the DNA methylation inhibitor to the patient may comprise administering imatinib mesylate to the patient for a period of time prior to the administration of the DNA methylation inhibitor, administering the DNA methylation inhibitor to the patient for a period of time prior to the administration of imatinib mesylate, or initiating administration of the DNA methylation inhibitor and imatinib mesylate to the patient at the same time. It is noted that the method may also comprise administering imatinib mesylate and the DNA methylation inhibitor to

the patient at the same time for at least a portion of the time that the drugs are administered.

According to any of the above methods, in one variation, imatinib mesylate is administered to the patient at a dose of 100-800 mg/day, optionally at a dose of 200-400 mg/day, and optionally at a dose of 500-800 mg/day. Such administrations may optionally last for a period of at least 2, 4, 6, 8, 10 or more days.

Also according to any of the above methods, in one variation, the DNA methylation inhibitor is administered to the patient via an intravenous infusion per day at a dose ranging from 1 to 100 mg/m², optionally at a dose ranging from 2 to 50 mg/m², and optionally at a dose ranging from 5 to 20 mg/m².

Also according to any of the above methods, the DNA methylation inhibitor may optionally be a cytidine analog such as cytosine arabinoside. In one variation, the cytidine analog is decitabine.

In one particular variation, the DNA methylation inhibitor is decitabine and is administered intravenously or subcutaneously. In a further particular variation, decitabine is administered to the patient via an intravenous infusion per day at a dose ranging from 1 to 100 mg/m², optionally ranging from 2 to 50 mg/m² and optionally ranging from 5 to 20 mg/m².

In one example, decitabine is administered to the patient via an intravenous infusion per day for at least 3 days per treatment cycle at a dose ranging from 1 to 100 mg/m². In a further example, decitabine is administered to the patient via an intravenous infusion at a dose ranging from 5 to 20 mg/m² for 1 hour per day for 5 consecutive days for 2 weeks per treatment cycle.

Compositions are also provided. In one embodiment, a composition is provided that comprises a DNA methylation inhibitor and imatinib mesylate. The DNA methylation inhibitor may optionally be a cytidine analog such as cytosine arabinoside. In one variation, the cytidine analog is decitabine. In another variation, the composition is formulated for intravenous, inhalation, oral, or subcutaneous administration.

DETAILED DESCRIPTION OF THE INVENTION

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185
10190
10195
10200
10205
10210
10215
10220
10225
10230
10235
10240
10245
10250
10255
10260
10265
10270
10275
10280
10285
10290
10295
10300
10305
10310
10315
10320
10325
10330
10335
10340

1. Imatinib Mesylate

Imatinib mesylate is a protein tyrosine kinase inhibitor that inhibits the Bcr-Abl tyrosine kinase created by the Philadelphia chromosome abnormality in CML. Imatinib mesylate achieves this inhibitory result through binding to the adenosine triphosphate-binding site of the Bcr-Abl tyrosine kinase, which prevents phosphorylation of substrates and related malignant transformation. Through inhibition of this kinase, it is believed that imatinib mesylate inhibits cell proliferation and induces apoptosis. T. Schindler et al (2000) Science 289:1938-1942.

Imatinib mesylate is indicated for treatment of CML patients in blast phase, accelerated phase, or in chronic phase after failure of interferon-alpha therapy. Present dosages recommended for treatment with imatinib mesylate are 400 mg/day for patients with chronic phase CML and 600 mg/day for patients with accelerated phase or blast phase CML. In the event of disease progression, failure to achieve a satisfactory hematologic response after at least 3 months of treatment; or loss of a previously achieved hematologic response, the dose of imatinib mesylate may be increased. Treatment dosage may be increased in patients with chronic phase CML from 400 mg/day to 600 mg/day in the absence of severe adverse drug reaction and severe non-leukemia related neutropenia or thrombocytopenia. Similarly, treatment dosage may be increased in patients with chronic phase CML from 600 mg/day to 800 mg/day. Novartis, Gleevec package insert T-2001-14 90012401.

However, many CML patients do not respond or lose response to treatment with imatinib mesylate. This is particularly the case in blast phase CML, the responses to imatinib mesylate are usually of very short duration, and most patients manifest resistant/refractory disease within six months of therapy. Druker et al (2001) N. Engl. J. Med. 344: 1038-1042. According to the present invention, it is believed that DNA methylation inhibitors can be used to treat CML patients that are resistant to the treatment with imatinib mesylate.

2. DNA Methylation Inhibitors

Decitabine, 5-aza-2'-deoxycytidine, is an antagonist of its related natural nucleoside, deoxycytidine. The only structural difference between these two compounds is the presence of a nitrogen at position 5 of the cytosine ring in decitabine as compared to a carbon at this position for deoxycytidine. Two isomeric forms of decitabine can be distinguished. The β -anomer is the active form. The modes of decomposition of decitabine in aqueous solution are (a) conversion of the active α -anomer to the inactive β -anomer (Pompon et al. (1987) J. Chromat. 388:113-122); (b) ring cleavage of the aza-pyrimidine ring to form N-(formylamido)-N'- β -D-2'-deoxy-(ribofuranosyl)-urea (Mojaverian and Repta (1984) J. Pharm. Pharmacol. 36:728-733); and (c) subsequent forming of guanidine compounds (Kissinger and Stemm (1986) J. Chromat. 353:309-318).

Decitabine possesses multiple pharmacological characteristics. At a molecular level, it is capable of specifically inhibiting cell growth at S phase and DNA methylation. At a cellular level, decitabine can induce cell differentiation and exert hematological toxicity. Despite having a short half life *in vivo*, decitabine has excellent tissue distribution.

The most prominent function of decitabine is its ability to specifically and potently inhibit DNA methylation. For example, in the methylation of cytosine in CpG islands, methylation of cytosine to 5-methylcytosine occurs at the level of DNA. Inside the cell, decitabine is first converted into its active form, the phosphorylated 5-aza-deoxycytidine, by deoxycytidine kinase which is primarily synthesized during the S phase of the cell cycle. The affinity of decitabine for the catalytic site of deoxycytidine kinase is similar to the natural substrate, deoxycytidine. Momparler et al. (1985) 30:287-299. After conversion to its triphosphate form by deoxycytidine kinase, decitabine is incorporated into replicating DNA at a rate similar to that of the natural substrate, dCTP. Bouchard and Momparler (1983) Mol. Pharmacol. 24:109-114.

Incorporation of decitabine into the DNA strand has a hypomethylation effect. Each class of differentiated cells has its own distinct methylation pattern.

After chromosomal duplication, in order to conserve this pattern of methylation, the 5-methylcytosine on the parental strand serves to direct methylation on the complementary daughter DNA strand. Substituting the carbon at the 5 position of the cytosine for a nitrogen interferes with this normal process of DNA methylation. The replacement of 5-methylcytosine with decitabine at a specific site of methylation produces an irreversible inactivation of DNA methyltransferase, presumably due to formation of a covalent bond between the enzyme and decitabine. Juttermann et al. (1994) Proc. Natl. Acad. Sci. USA 91:11797-11801. By specifically inhibiting DNA methyltransferase, the enzyme required for methylation, the aberrant methylation of the tumor suppressor genes can be prevented.

3. Treatment of Different Disease States

Described herein are several different disease states which may be treated by the combination therapy methods and compositions provided herein. It is noted that other disease states that may be treated with imatinib mesylate may also be treated with the combination of imatinib mesylate and a DNA methylation inhibitor where the DNA methylation inhibitor synergistically renders the imatinib mesylate more effective, for example by reducing resistance to imatinib mesylate that the patient may naturally have or may develop over time.

A. Chronic myelogenous leukemia

Unlike other forms of leukemia, CML is a homogeneous disease. Almost all patients with CML (90%) have the same chromosomal abnormality in their leukemic cells. In up to 40% of CML patients, the disease progresses directly from the chronic to the blastic phase. Cortes and Kantarjian (1998) in Cancer management: a multidisciplinary approach, 2d ed. Huntington, NY: Publisher Research Management, Inc, p306-15. Blast transformation typically occurs at 3 to 5 years, but onset is random and may be observed at the time of initial diagnosis. Common features for chronic phase CML patients include fatigue, weight loss, and signs or symptoms of splenomegaly (e.g., left upper quadrant pain, abdominal fullness, a

palpable mass), which occurs in up to 70% of patients. Cortes et al (1996) *Am J Med* 100(5):555-70. Patients with extreme elevations in white blood cell count (hyperleukocytosis) may demonstrate signs or symptoms of leukostasis, including mental status changes, focal neurologic deficits, tinnitus, dyspnea, and priapism. Retinal hemorrhages or other bleeding manifestations may appear irrespective of platelet counts. This is attributable to the qualitative platelet dysfunction that is common in CML. Goldman (1997) *BMJ (Clin Res Ed)* 314:657-60.

As CML progresses from chronic phase to accelerated phase, the patient may experience fever, night sweats, weight loss, and progressive splenomegaly. More often, however, there is no significant change in symptoms, and onset is heralded by hematologic progression (e.g., worsening of blood counts) or cytogenetic evolution (e.g., development of new chromosomal abnormalities). As CML progresses from the accelerated phase to blast phase, the patient may experience fever, weight loss, night sweats, bone pain, and constitutional symptoms. Lymphadenopathy, leukemia cutis, central nervous system disease, and bleeding secondary to progressive thrombocytopenia also may occur. Hughes and Goldman (1995) in *Hematology: basic principles and practice*, 2d ed. New York: Churchill Livingstone, p1142-59.

CML staging involves the use of a system of categorization to describe the seriousness of the CML and attempts to put patients with similar prognosis and treatment in the same staging group. Staging is important in developing CML treatment strategies as it allows doctors to compare the efficacy of different treatments for patients with similar conditions, and aids in the determination of treatment decisions. Staging systems for CML are generally based on clinical features with demonstrated prognostic significance, including age, size of spleen, percentage of circulating and marrow blasts, degree of basophilia, extent of thrombocytosis, and presence of atypical chromosomal abnormalities. Faderl, S., Talpaz, M., Estrov, Z., and Kantarjian, H. M. Chronic myelogenous leukemia: biology and therapy. *Ann. Intern. Med.*, 131: 207-219, 1999; Grier HE, Civin CL. Myeloid Leukemias, Myelodysplasia, and Myeloproliferative Diseases in Children. In (Nathan and Oski, eds) *Hematology of Infancy and Childhood*, volume 2nd. 5th

10
5
Edition, W. B. Saunders Company, 1998; 34:1286-1459; Santini, V., Kantarjian, H. M., and Issa, J. P. Changes in DNA methylation in neoplasia: pathophysiology and therapeutic implications. *Ann. Intern. Med.*, 134: 573-586, 2001; Lubbert M, Wijermans P, Kunzmann R, et al. Cytogenetic responses in high-risk myelodysplastic syndrome following low-dose treatment with the DNA methylation inhibitor 5-aza-2'-deoxycytidine. *Br J Haematol* 2001 Aug; 114(2):349-357. In addition, specific response to interferon alpha-2b therapy is considered to be a particularly sensitive predictor of long-term survival following treatment. Goldman (1997) *Baillieres Clin Haematol* 10(2):405-21.

10
15
20
Prior to the administration of a DNA methylation inhibitor, CML patients may optionally be staged to determine the severity of the CML. Staging may include determining the number of blasts, promyelocytes, basophil, and platelets per liter of peripheral blood or bone marrow.

10
15
20
25
Patients with CML in the chronic phase have all of the following conditions: less than 15% blasts in the peripheral blood or bone marrow; less than 30% blasts and promyelocytes in the peripheral blood or bone marrow; less than 20% basophils in the peripheral blood; 100 times 10 supra 9 per liter platelets; and no extramedullary involvement other than liver or spleen. Patients with CML in the accelerated phase have at least one of the following conditions: 15% to less than 30% blasts in the peripheral blood or bone marrow; 30% blasts and promyelocytes in the peripheral blood or bone marrow (but less than 30% blasts in the peripheral blood or bone marrow); 20% basophils in the peripheral blood; or less than 100 times 10 supra 9 per liter platelets.

25
30
Blast phase CML patients have at least 30% blasts and promyelocytes in the peripheral blood or bone marrow, or extramedullary involvement other than liver or spleen.

30
Preferable blast phase CML patients treated using the compositions and methods of the present invention are two years of age or older, have histologically confirmed diagnosis of blast phase CML, were previously treated with imatinib mesylate with no response or resulted in loss of response, and have bilirubin levels

greater than or equal to 1.5 times the upper limit of normal, serum glutamic-oxaloacetic transaminase and serum glutamic-pyruvic transaminase levels greater than or equal to 2.5 times the upper limit of normal, and serum creatine levels greater than or equal to 1.5 times the upper limit of normal.

5 Patients with CML in chronic, accelerated or blast phase, who are resistant to standard therapy generally have a relatively poor prognosis with a rapid progression to blast crisis, bone marrow failure and death. To measure the success of a therapeutic treatment of CML, hematologic responses are determined. A complete hematologic response is defined as maintaining the following conditions for four weeks: less than or equal to 5% blasts in bone marrow; no peripheral blood blasts; absolute neutrophil count of greater than 1.5 times 10×10^9 per liter; platelet count of greater than 100 times 10^9 per liter; and no extramedullary involvement.

10
10071849-620702
18
20 The Philadelphia chromosome, with the Bcr-Abl oncogene detectable at the molecular level, is present at diagnosis in 95% of patients. Optionally, complete hematologic responses are further classified according to suppression of the Philadelphia chromosome (Ph). For example, a patient with complete hematologic response and greater than 65% Ph positive is classified as no cytogenetic response. Patient responses that are 36% to 65% Ph positive are classified as minimal cytogenetic responses, 1% to 35% Ph positive are partial cytogenetic responses, and 0% Ph positive are complete cytogenetic responses.

20 A partial hematologic response is defined as maintaining the following conditions for four weeks: less than or equal to 5% blasts in bone marrow; no peripheral blood blasts; absolute neutrophil count of less than 1.5 times 10×10^9 per liter; and platelet count of less than 100 times 10^9 per liter.

25 A hematologic improvement is defined as maintaining the following conditions for four weeks: less than 15% blasts in bone marrow and peripheral blood; less than 40% blasts and promyelocytes in peripheral blood and bone marrow; less than 20% basophils in peripheral blood; and no extramedullary involvement other than liver or spleen.

CML disease progression is associated with hypermethylation of a promoter region within Bcr-Abl. J-P Issa et al (1999) Blood 93:2075-2080. In mammalian cells, approximately 3% to 5% of the cytosine residues in genomic DNA are present as 5-methylcytosine. Ehrlich et al (1982) Nucleic Acid Res. 10:2709-2721. This modification of cytosine takes place after DNA replication and is catalyzed by DNA methyltransferase using S-adenosyl-methionine as the methyl donor. Approximately 70% to 80% of 5-methylcytosine residues are found in the CpG sequence. Bird (1986) Nature 321:209-213. This sequence, when found at a high frequency, in the genome, is referred to as CpG islands. Unmethylated CpG islands are associated with housekeeping genes, while the islands of many tissue-specific genes are methylated, except in the tissue where they are expressed. Yevin and Razin (1993) in DNA Methylation: Molecular Biology and Biological Significance. Basel: Birkhauser Verlag, p523-568. This methylation of DNA has been proposed to play an important role in the control of expression of different genes in eukaryotic cells during embryonic development. Consistent with this hypothesis, inhibition of DNA methylation has been found to induce differentiation in mammalian cells. Jones and Taylor (1980) Cell 20:85-93.

Methylation of DNA in the regulatory region of a gene can inhibit transcription of the gene. This may be because 5-methylcytosine protrudes into the major groove of the DNA helix, which interferes with the binding of transcription factors.

The methylated cytosine in DNA, 5-methylcytosine, can undergo spontaneous deamination to form thymine at a rate much higher than the deamination of cytosine to uracil. Shen et al. (1994) Nucleic Acid Res. 22:972-976. If the deamination of 5-methylcytosine is unrepaired, it will result in a C to T transition mutation. For example, many "hot spots" of DNA damages in the human p53 gene are associated with CpG to TpG transition mutations. Denissenko et al. (1997) Proc. Natl. Acad. Sci. USA 94:1893-1898.

B. Treatment of Chronic Myelogenous Leukemia

In a particular embodiment, a method is provide for treating CML. It has been found that unchecked production of tyrosine kinase Bcr-Abl leads to excessive levels of white blood cells in the blood and bone marrow, but disrupts the normal production of white blood cells; and imatinib mesylate works specifically to block the activity of Bcr-Abl tyrosine kinase. However, methylation of Bcr-Abl has been found to correlate with progression with CML: hypermethylation in 24-68% of CML patients in the chronic phase, while patients in the accelerated phase and blast crisis had hypermethylation frequencies at 73% and 80%, respectively. Issa et al. (1999) Blood 93:2075-2080.

Other than Bcr-Abl gene, multiple genes, including the p15/Ink-4b cell-cycle regulator gene, are found to contain aberrant methylation in more advanced stages of CML. Cortes et al (1997) Baillieres Clin Haematol 10(2):277-90. Aberrations in DNA methylation, whether general or site specific, are common in cancer and have important roles in tumor initiation, progression and resistance. Lubbert et al (2001) Br J Haematol 114(2):349-357. Methylation of the p15 promoters is associated with progression of CML. Nguyen et al. (2000) Blood 95:2990-2992.

The inventor believes that hypermethylation of p15 would lead to loss of its tumor suppression function and accelerates transformation of Bcr-Abl, ultimately leading to resistance to the treatment of CML with imatinib mesylate. Administering a DNA methylation inhibit to a CML patient may re-establish the tumor suppression functions of p15, which, in turn, leads to sensitization of the patient to the treatment with imatinib mesylate.

The present invention provides a method of treating a patient having chronic myelogenous leukemia, comprising: administering to a patient having chronic myelogenous leukemia but resistant to the treatment with imatinib mesylate a therapeutically effective amount of a DNA methylation inhibitor. In one variation, the patient has already manifested resistance to imatinib mesylate within 6 months of the treatment with imatinib mesylate as defined by no improvement in the prognosis or worsening of the prognosis.

10
10071849-020702
The DNA methylation inhibitor may be optionally administered to the patient in the blast, chronic or accelerated phase of chronic myelogenous leukemia. In one embodiment of the present invention, a method is provided for treating patients having blast phase CML, where prior to treatment, the patient's CML is staged. The method comprises administering to a patient suffering from blast phase CML, after the CML has been staged, a DNA methylation inhibitor. Optimally, staging the CML includes determining the number of blasts, promyelocytes, basophil, and platelets per liter of peripheral blood or bone marrow. Optionally, the patient in blast phase of CML has more than 30% blasts in the peripheral blood or bone marrow.

1
A method is also provided for treating a patient having chronic myelogenous leukemia but manifesting intolerance to imatinib mesylate. The method comprises: administering to the patient a therapeutically effective amount of a DNA methylation inhibitor which mitigates the imatinib mesylate intolerance.

20
Preferably the DNA methylation inhibitor is administered to a patient who has already manifested intolerance to imatinib mesylate within 6 months of the treatment with imatinib mesylate. The patient's intolerance to imatinib mesylate can be defined by manifesting symptoms or adverse effects such hepatotoxicity, fluid retention syndrome, neutropenia, hemorrhage, dyspepsia, dyspnea, diarrhea, muscle cramps, skin rash, fatigue, headache, nausea, vomiting, and thrombocytopenia.

A method is also provided for treating a patient having chronic myelogenous leukemia, comprising: co-administering to the patient imatinib mesylate and decitabine such that the patient's resistance to imatinib mesylate in the absence of decitabine is reduced.

25
In one variation, imatinib mesylate is administered to the patient at a dose of 100-800 mg/day. For chronic phase CML patients, imatinib mesylate is preferentially administered at a dose of 200-400 mg/day. For accelerated or blast phase CML patients, imatinib mesylate is preferably administered at a dose of 500-800 mg/day.

5

15

20

24

10071849-020702
5 The inventor believes that while imatinib mesylate can also inhibit tyrosine kinase activity of PDGF-R, as in CML loss of p15 tumor suppression function may lead to transformation of PDGF-R, ultimately leading to resistance to imatinib mesylate treatment. By using the composition and methods of the present invention, diseases associated with tyrosine kinase activity of PDGF-R can be treated more efficaciously and with significantly reduced adverse effects.

15 The compositions and methods of the present invention may also be used to treat diseases associated with tyrosine kinase activity of the transmembrane protein c-Kit such as gastrointestinal stromal tumor (GIST) and small cell lung cancer (SCLC). c-Kit is defined by the CD117 antigen and is the product of the *c-kit* proto-oncogene. Activating or gain-of-function mutations in the *c-kit* gene have been identified in the majority of GISTs. As a result, Kit is constitutively expressed to provide growth and survival signals to GIST cells, which are crucial to the pathogenesis of the disease. So far, GIST has been found to be the most common mesenchymal tumor of the GI tract and resistant to chemotherapy and radiation treatment. Recently, the US Food and Drug Administration (FDA) has approved imatinib mesylate (or GLEEVEC®) for the treatment of patient with KIT (or CD117)-positive unresectable and/or metastatic magligant GISTs.

20 The inventor believes that while imatinib mesylate can suppress GIST growth by inhibiting the tyrosine kinase activity of c-Kit, mutations in *c-kit* may render the disease less responsive to the treatment with imatinib mesylate. In addition, hypermethylation in the genes participating in the signal transduction pathway of c-Kit would also lead to transformation of c-Kit, ultimately leading to resistance to imatinib mesylate treatment. By targeting c-Kit and DNA
25 hypermethylation using the combination therapy of the present invention, diseases associated with tyrosine kinase activity of c-Kit can be treated more efficaciously and with significantly reduced adverse effects.

4. Routes of Delivery, Formulations and Kits

The DNA methylation inhibitor employed in the present invention may be administered or coadministered in any conventional dosage form. For example, the inhibitor be administered or coadministered parenterally, orally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery by catheter or stent, subcutaneously, intraadiposally, intraarticularly, or intrathecally.

In a preferred embodiment, decitabine is administered to a patient by injection, including intravenous or subcutaneous injection, such as bolus intravenous injection, continuous intravenous infusion and intravenous infusion over 1 hour. For example, decitabine may administered into the patient via an 1-24 hour intravenous infusion per day for 3-5 days per treatment cycle at a dose preferably ranging from 1-100 mg/m², more preferably ranging from 2-50 mg/m², and most preferably from 5-20 mg/m². The preferred dosage below 50 mg/m² for decitabine is considered to be much lower than that used in conventional chemotherapy for cancer.

In another embodiment, decitabine is administered via intravenous infusion at a dose ranging from 1 to 100 mg/m² per day for at least 3 days per treatment cycle. In yet another embodiment, decitabine is administered via intravenous infusion at a dose ranging from 5 to 20 mg/m² for 1 hour per day for 5 consecutive days for 2 weeks per treatment cycle.

The DNA methylation inhibitors employed in the invention may also be administered or coadministered in slow release dosage forms. Furthermore, the DNA methylation inhibitors may be administered or coadministered with conventional pharmaceutical excipients and additives.

Decitabine may be supplied as sterile powder for injection, together with buffering salt such as potassium dihydrogen and pH modifier such as sodium hydroxide. This formulation is preferably stored at 2-8°C, which should keep the drug stable for at least 2 years. This powder formulation may be reconstituted with

10 ml of sterile water for injection. This solution may be further diluted with infusion fluid known in the art, such as 0.9% sodium chloride injection, 5% dextrose injection and lactated ringer's injection. It is preferred that the reconstituted and diluted solutions be used within 4-6 hours for delivery of maximum potency.

5 The inventive combination of DNA methylation inhibitor and imatinib mesylate may be used in the form of kits. The arrangement and construction of such kits is conventionally known to one of skill in the art. Such kits may include containers for containing the inventive combination of therapeutic agents and/or compositions, and/or other apparatus for administering the inventive combination of therapeutic agents and/or compositions.

EXAMPLES

Example 1:

10 Patients with CML receive decitabine at 15 mg/m² IV over one hour on five consecutive days each week (e.g., Monday through Friday) for two weeks, followed by four weeks rest. For patients who achieve and remain in remission, treatment may continue so long as no adverse drug experiences occur and there is no evidence of disease progression or relapse.

15 Since the effect of decitabine may be delayed for up to two cycles, patients with high white blood cell counts ($>10 \times 10^9$ per liter) at study entry may optionally receive hydroxyurea (HU) 1-3 g orally daily concomitantly with the first two cycles of decitabine only. HU may be continued until the end of dosing in the second cycle of decitabine (e.g., the 13th day of the second cycle) or until the white blood cell count falls below 10×10^9 per liter during the second cycle. This is done to minimize the risks of complications due to hyperleukocytosis while awaiting a response. In general, HU can be given a 1 g daily if the white blood cell count is between $10-40 \times 10^9$ per liter, 2 g daily if the white blood cell count is $40-50 \times 10^9$ per liter, and 3 g daily if the white blood cell count is above 50×10^9 per liter. Other dose schedules may be used as indicated. Children with white blood cell counts above $10-20 \times 10^9$ per liter can be treated with HU at 10-20 mg/kg/day. The use of anagrelide is

allowed for platelets $>600 \times 10^9$ per liter, but is generally discontinued prior to decitabine dosing in the third cycle.

Patients achieving an unconfirmed complete or partial hematological response, or a hematologic improvement, will continue to receive therapy every six weeks until they show evidence of disease progression or relapse.

The second and subsequent cycles of decitabine will be instituted when patients have recovered from hematologic or non-hematological toxicity. Hematologic recovery is generally an absolute neutrophil count $>1 \times 10^9$ per liter or to baseline and platelets to $>150 \times 10^9$ per liter, untransfused for the preceding week.

The decitabine dose may be escalated incrementally by 25% to reduce the absolute neutrophil count to between 1×10^9 per liter and 1.5×10^9 per liter and a platelet count to between 50×10^9 per liter and 100×10^9 per liter with each cycle. The decitabine dose may be decreased incrementally by 25% in the face of grade 4 hematologic toxicity or grade 3 or 4 non-hematological toxicity, other than alopecia.

It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this invention provided they come within the scope of the appended claims and their equivalents. Additionally, the above examples are provided for the purpose of illustrating the claimed invention, and should not be construed so as to limit the scope of the claimed invention.